ELSEVIER

Carbohydrate Research 262 (1994) C7–C9

CARBOHYDRATE RESEARCH

CONTENTS

1,3-Parallel C//O interactions in acyclic carbohydrates: the crystal and molecular structure of 1-deoxy-1-nitro-D-altritol
P. Köll, M. Bischoff (Oldenburg, Germany), C. Bretzke and J. Kopf (Hamburg, Germany) . . 1

1,3-Parallel O//O interactions in acyclic carbohydrates: the crystal and molecular structures of *meso*-D-*glycero*-L-*altro*-heptitol and the monohydrate, *meso*-D-*glycero*-L-*ido*-heptitol, and *meso*-D-*glycero*-L-*altro*-heptitol and D-*glycero*-L-*galacto*-heptitol heptaacetates
J. Kopf, M. Morf, B. Hagen (Hamburg, Germany), M. Bischoff and P. Köll (Oldenburg, Germany) 9

Cellulose char structure: a combined analytical Py–GC–MS, FTIR, and NMR study
I. Pastorova (Amsterdam, Netherlands), R.E. Botto, P.W. Arisz (Argonne, IL, USA) and J.J. Boon (Amsterdam, Netherlands) 27

Benzyl 7-*O*-allyl-2,3-*O*-isopropylidene-4-*O*-*p*-methoxybenzyl-L-*glycero*-α-D-*manno*-heptopyranoside, a versatile LD-manHepp derivative
B. Grzeszczyk and A. Zamojski (Warsaw, Poland) 49

The preparation of intermediates for the synthesis of 1D-*myo*-inositol 1,4,5- and 2,4,5-trisphosphates, 1,4-bisphosphate 5-phosphorothioate, and 4,5-bisphosphate 1-phosphorothioate from 1D-3,6-di-*O*-benzyl-1,2-*O*-isopropylidene-*myo*-inositol
T. Desai, J. Gigg, R. Gigg and E. Martín-Zamora (London, United Kingdom) 59

Synthesis of double-chain bis-sulfone neoglycolipids of the 2″-, 3″-, 4″-, and 6″-deoxyglobotrioses
Z. Zhiyuan and G. Magnusson (Lund, Sweden) 79

Synthesis of a 2,3-dideoxy-2,3-difluorofuranose with the D-*lyxo* configuration. An intramolecular rearrangement of methyl 5-*O*-benzoyl-2,3-dideoxy-2,3-difluoro-D-lyxofuranoside observed during the attempted synthesis of 1-(2,3-dideoxy-2,3-difluoro-β-D-lyxofuranosyl)thymine
L.S. Jeong, B.B. Lim and V.E. Marquez (Bethesda, MD, USA) 103

Chemical characteristics of insoluble glucans from the cell wall of the marine green alga *Ulva lactuca* (L.) Thuret
M. Lahaye, D. Jegou and A. Buleon (Nantes, France) 115

Chemical methylation of agaroid hydroxyl-groups
I.J. Miller, R. Falshaw and R.H. Furneaux (Lower Hutt, New Zealand) 127

Notes

Synthesis of 6^5- and/or 4^5-*O*-glycosyl *p*-nitrophenyl α-maltopentaoside derivatives for the differential assay of human α-amylases
K. Ozawa (Gifu, Japan), Y. Yamagata, S. Satomura, H. Ishida (Hyogo, Japan), M. Kiso and A. Hasegawa (Gifu, Japan). 137

Stereochemical control in the formation of thiazolidines from O-protected reducing sugars
A. Wadouachi, D. Beaupere, R. Uzan, I. Stasik (Amiens, France), D.F. Ewing and G. Mackenzie (Hull, United Kingdom) 147

Structural characterization of alkali-soluble glucans from the mycelium of *Aspergillus giganteus* mut. *alba*
P. Rouhier, M. Bruneteau, G. Michel (Villeurbanne, France), J. Fiema and A. Zurzycka (Cracow, Poland) 155

Molecular mass distribution of chitin and chitosan
M. Hasegawa, A. Isogai and F. Onabe (Tokyo, Japan) 161

Detection of cellulose in the cell wall of some red algae by ^{13}C NMR spectroscopy
R. Toffanin, S.H. Knutsen, C. Bertocchi, R. Rizzo and E. Murano (Trieste, Italy) 167

Regular papers

Studies of the neutral trisaccharides of goat (*Capra hircus*) colostrum and of the one- and two-dimensional ^{1}H and ^{13}C NMR spectra of 6′-*N*-acetylglucosaminyllactose
T. Urashima, W.A. Bubb, M. Messer (Sydney, NSW, Australia), Y. Tsuji and Y. Taneda (Hokkaido, Japan) 173

A ^{13}C solid-state NMR study of ion-exchange resins derived from natural polysaccharides
K.R. Morgan, J.A. Hemmingson, R.H. Furneaux and R.A. Stanley (Lower Hutt, New Zealand) 185

Synthesis of the 5-aminopentyl glycoside of β-D-Gal*p*-(1 → 4)-β-D-Glc*p*NAc-(1 → 3)-L-Fuc*p* and fragments thereof related to glycopeptides of human Christmas factor and the marine sponge *Microciona prolifera*
T. Ziegler (Stuttgart, Germany) 195

The reaction between epichlorohydrin and polysaccharides: Part 1, syntheses of some model substances with non-cyclic substituents
L. Holmberg (Uppsala, Sweden), B. Lindberg (Stockholm, Sweden) and B. Lindqvist (Uppsala, Sweden) 213

Synthesis of carboxyl-reduced analogues related to the *Chlamydia*-specific Kdo trisaccharide epitope
F.W. D'Souza, P. Kosma (Wien, Austria) and H. Brade (Borstel, FRG) 223

α-(1 → 2)-, α-(1 → 3)-, and α-(1 → 6)-Linked thioglycosidic disaccharides: syntheses and anti-HIV testing of thiokojibiose octaacetate, thionigerose, and thioisomaltose
R.N. Comber, J.D. Friedrich, D.A. Dunshee, S.L. Petty and J.A. Secrist III (Birmingham, AL, USA) 245

The preparation and characterization of some Amadori compounds (1-amino-1-deoxy-D-fructose derivatives) derived from a series of aliphatic ω-amino acids
V.V. Mossine, G.V. Glinsky and M.S. Feather (Columbia, MO, USA) 257

Selective pivaloylation and diphenylacetylation of cyclomalto-oligosaccharides
F. Santoyo-González, J. Isac-García, A. Vargas-Berenguel, R. Robles-Díaz and F.G. Calvo-Flores (Granada, Spain) 271

Synthesis of a di-*O*-methylated pentasaccharide for use in the assay of *N*-acetylglucosaminyltransferase III activity
S.H. Khan, C.A. Compston, M.M. Palcic and O. Hindsgaul (Edmonton, AB, Canada) 283

Synthesis of transition-state analogues as potential inhibitors of sialidase from *Influenza* virus
P.-A. Driguez (Villeurbanne, France), B. Barrere, G. Quash (Oullins, France) and A. Doutheau (Villeurbanne, France) 297

Structural investigation of the capsular polysaccharide of *Escherichia coli* O101 : K103 : H^- using bacteriophage degradation and NMR spectroscopy
M.R. Grue, H. Parolis and L.A.S. Parolis (Grahamstown, South Africa) 311

Structural elucidation of the capsular polysaccharide produced by *Escherichia coli* O20 : K84 : H26
D.V. Whittaker, L.A.S. Parolis and H. Parolis (Grahamstown, South Africa) 323

Synthesis of 2-deoxy-α-D-*arabino*-hexopyranosyl phosphate and 2-deoxy-maltooligosaccharides with phosphorylase
B. Evers, P. Mischnick and J. Thiem (Hamburg, Germany) 335

Author index C1

Subject index C3

Contents C7